


§4.1 Rational Functions and Asymptotes


Rational Function - a function of the form $f(x) = \frac{p(x)}{q(x)}$
where $p(x)$ and $q(x)$ are polynomials with $q(x) \neq 0$.

The graphs of rational functions approach (get closer and closer to) lines called **asymptotes**.

Basic Graphs (memorize these)



$$y = \frac{1}{x}$$




$$y = \frac{1}{x^2}$$

Example 1 Use stretching/shrinking, reflecting and shifting rules to graph the following.

a.) $f(x) = \frac{1}{x-2} + 1$

b.) $f(x) = -\frac{2}{(x+4)^2}$



To Find the Asymptotes of a Rational Function:

- (1) Vertical Asymptotes - Find any vertical asymptotes by setting the denominator equal to 0 and solving for x to get the equation

$$\boxed{x = a} .$$

- (2) Horizontal Asymptotes

Rule 1 If the numerator has lower degree than the denominator, the horizontal asymptote is $\boxed{y = 0}$.

Rule 2 If the numerator and denominator have the same degree and a_n is the leading coefficient of the numerator and b_n is the leading coefficient of the denominator, the horizontal asymptote is

$$\boxed{y = \frac{a_n}{b_n}} .$$

Rule 3 If the numerator has higher degree than the denominator, there is $\boxed{\text{no horizontal asymptote}}$.

- (3) Slant Asymptotes - If the numerator is of degree exactly one more than the denominator, there is an slant asymptote. To find it, divide the numerator by the denominator and disregard any remainder. The equation of the slant asymptote is the result of setting y equal to the quotient.

Example 2 Give the equations of the vertical, horizontal and/or slant asymptotes of the rational function.

a.) $f(x) = \frac{3x}{(x+1)(x-2)}$

b.) $f(x) = \frac{(x-5)(x-2)}{x^2+9}$

c.) $f(x) = \frac{2x^2+3}{x-4}$

d.) $f(x) = \frac{2(3x-1)(x+4)}{(x+2)(5x-3)}$

Example 3 Find the x-intercepts and y-intercept of the rational function.

a.) $f(x) = \frac{3x}{(x+1)(x-2)}$

b.) $f(x) = \frac{(x-5)(x-2)}{x^2+9}$