

## §7.3 Trigonometric Identities

### Fundamental Trigonometric Identities

#### Reciprocal Identities

$$\sin \theta = \frac{1}{\csc \theta} \quad \cos \theta = \frac{1}{\sec \theta} \quad \tan \theta = \frac{1}{\cot \theta}$$

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

#### Quotient or Ratio Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

#### Pythagorean Identities

$$\sin^2 \theta + \cos^2 \theta = 1 \quad \tan^2 \theta + 1 = \sec^2 \theta \quad 1 + \cot^2 \theta = \csc^2 \theta$$

#### Cofunction Identities

$$\begin{array}{lll} \sin(90^\circ - \theta) = \cos \theta & \cos(90^\circ - \theta) = \sin \theta & \tan(90^\circ - \theta) = \cot \theta \\ \csc(90^\circ - \theta) = \sec \theta & \sec(90^\circ - \theta) = \csc \theta & \cot(90^\circ - \theta) = \tan \theta \end{array}$$

#### Even and Odd Trigonometric Functions

The cosine and secant functions are **even**.

$$\cos(-t) = \cos t \quad \sec(-t) = \sec t$$

The sine, cosecant, tangent, and cotangent functions are **odd**.

$$\sin(-t) = -\sin(t) \quad \csc(-t) = -\csc(t)$$

$$\tan(-t) = -\tan(t) \quad \cot(-t) = -\cot(t)$$

Example 1      Simplify :  $\frac{\cot \theta}{\csc \theta}$

Example 2      Simplify:  $\frac{\cos \theta}{1 + \sin \theta}$

Example 3      Simplify:  $\frac{1 + \sin u}{\sin u} + \frac{\cot u - \cos u}{\cos u}$

Example 4      Simplify:  $\frac{\sin^2 \theta - 1}{\tan \theta \sin \theta - \tan \theta}$

## **Establish (prove) Trigonometric Identities**

- an identity is an equation which is true for all values for which the equation is defined
- to verify an identity, generally work with one side of the equation and show that it equals the other side
- some suggestions to consider when verifying identities:

- 1.) simplify the more complex side
- 2.) perform algebraic operations including squaring, factoring, adding or subtracting fractions, multiplying the numerator and denominator by a nonzero factor
- 3.) rewrite in terms of sine and cosine
- 4.) rewrite in terms of a single trigonometric function
- 5.) use other identities (reciprocal identities, ratio identities, Pythagorean identities)

Examples Verify each identity.

a.)  $\csc \theta \cdot \tan \theta = \sec \theta$

$$\text{b.) } \sin^2(-\theta) + \cos^2(-\theta) = 1$$

$$\text{c.) } \frac{\sin^2(-\theta) - \cos^2(-\theta)}{\sin(-\theta) - \cos(-\theta)} = \cos \theta - \sin \theta$$

$$d.) \frac{1 + \tan u}{1 + \cot u} = \tan u$$

$$e.) \frac{\sin \theta}{1 + \cos \theta} + \frac{1 + \cos \theta}{\sin \theta} = 2 \csc \theta$$

$$f.) \frac{\tan v + \cot v}{\sec v \csc v} = 1$$